



Calculation of Equation of State of QCD at Zero Temperature and Finite Chemical Potential

Hong-Shi Zong^{a,b} Yu Jiang^a Xiao-ya Li^c Wei-Min Sun^{a,b}

^a Department of Physics, Nanjing University

^bJoint Center of Particle, Nuclear Physics and Cosmology, Nanjing University-Purple
Mountain Observatory

^c Department of Physics, Sichuan University

[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 1 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



Outline

1. Introduction
2. Formula for EOS of QCD at Zero Temperature and Finite Chemical Potential
3. Dyson-Schwinger Equation (DSE) of QCD
4. Calculation of EOS with Rainbow-ladder DSE
5. Calculation of EOS with Hard-Dense-Loop Approximation
6. Calculation of EOS with Quasi-particle Model
7. Calculation of EOS with a Nonperturbative Approach Inspired by Chiral Perturbation Theory
8. Summary

[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 2 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



List of part of published papers during 2008.1-2010.6:

1. H. T. Feng, W. M. Sun, D. K. He, and H. S. Zong, *Influence of Finite Chemical Potential on the Fermion Chiral Condensate in QED₃*, Phys. Lett. B. **661** (2008), 57.
2. M. He, Y. Jiang, W. M. Sun, and H. S. Zong, *Chiral Susceptibility in an Effective Interaction Model*, Phys. Rev. **D77**, 076008 (2008).
3. Y. Jiang, Y. B. Zhang, W. M. Sun, and H. S. Zong, *Response of Quark Condensate to the Chemical Potential*, Phys. Rev. **D78**, 014005 (2008).
4. Y. Jiang, Y. M. Shi, H. T. Feng, W. M. Sun, H. S. Zong, *Quark-meson vertices and pion properties at finite chemical potential*, Phys. Rev. **C78**, 025214 (2008).
5. H. S. Zong and W. M. Sun, *The Calculation of the Equation of State of QCD at Finite Chemical Potential and Zero Temperature*, Phys. Rev. **D78**, 054001 (2008).
6. L. Chang, Y. X. Liu, W. M. Sun, and H. S. Zong, *Revisiting the Vector and Axial Vector Vacuum Susceptibilities*, Phys. Lett. **B669**, 327 (2008).

[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 3 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



7. Y. Jiang, Y. M. Shi, H. Li, W. M. Sun, H. S. Zong, *Calculation of f_π and m_π at finite chemical potential*, Phys. Rev. **D78**, 116005 (2008).
8. M. He, J. F. Li, W. M. Sun, and H. S. Zong, *Quark Number Susceptibility Around the Critical End Point*, Phys. Rev. **D79**, 036001 (2009).
9. L. Chang, Y. X. Liu, C. D. Roberts, Y. M. Shi, W. M. Sun, and H. S. Zong, *Chiral Susceptibility and the Scalar Ward Identity*, Phys. Rev. **C79**, 035209 (2009).
10. M. He, F. Hu, W. M. Sun, and H. S. Zong, *Crossover from a continuum study of chiral susceptibility*, Phys. Lett. **B675**, 32 (2009).
11. D. K. He, X. X. Ruan, Y. Jiang, W. M. Sun, and H. S. Zong, *The model study of quark-number susceptibility at finite chemical potential and temperature*, Phys. Lett. **B680**, 432 (2009).
12. X. Y. Li, X. F. Lü, B. Wang, W. M. Sun, and H. S. Zong, *Study of the properties of cold dense nuclear matter based on a non-perturbative approach inspired by chiral perturbation theory*, Phys. Rev. **C80**, 034909 (2009).
13. L. Chang, Y. X. Liu, C. D. Roberts, Y. M. Shi, W. M. Sun, and H. S. Zong, *Vauum pseudoscalar susceptibility*, Phys. Rev. **C81**, 032201(R) (2010).

访问主页

标题页

◀ ▶

◀ ▶

第 4 页 共 49 页

返 回

全屏显示

关 闭

退 出



14. Y. Jiang, H. X. Zhu, W. M. Sun, and H. S. Zong, *The quark number susceptibility at Hard-Thermal-Loop approximation*, J. Phys. **G 37**, 055001 (2010).
15. H. T. Feng, M. He, W. M. Sun, and H. S. Zong, *Investigation of Phase Transition in QED₃*, Phys. Lett. **B688**, 178 (2010).
16. Y. Jiang, N. Li, W. M. Sun, and H. S. Zong, *The calculation of quark-number susceptibility at finite chemical potential and temperature*, Nucl. Phys. **A834**, 587C (2010).
17. H. B. Zeng, Z. Y. Zhe, and H. S. Zong, *Superconducting coherence length and magnetic penetration depth of a p-wave holographic superconductor*, Phys. Rev. **D81**, 106001 (2010).
18. J. F. Li, S. S. Huang, H. T. Feng, W. M. Sun and H. S. Zong, *Dynamical symmetry breaking in QED₃ with Abelian Higgs model*, Symmetry. **2**, 907 (2010).
19. Y. Jiang, H. Li, S. S. Huang, W. M. Sun, and H. S. Zong, *The equation of state and quark number susceptibility in Hard-Dense-Loop approximation*, to be published in J. Phys. **G**.

访问主页

标 题 页

◀ ▶

◀ ▶

第 5 页 共 49 页

返 回

全 屏 显 示

关 闭

退 出



★ Introduction

The QCD thermodynamical system at finite temperature (T) and finite chemical potential (μ) is described by the density matrix

$$\hat{\rho} = e^{-(\hat{H}_{QCD} - \mu \hat{N}_B)/T}$$

Partition function: $\mathcal{Z}[\mu, T] = \text{Tr} \hat{\rho} = \sum_{\alpha} \exp\left\{-\frac{E_{\alpha} - \mu N_{\alpha}}{T}\right\}$

1. The partition function plays an important role in thermal field theory. From the partition function one can obtain all thermodynamical quantities and the equation of state (EOS).
2. A neutron star can be approximately regarded as a zero temperature and high density system. The QCD partition function and EOS at zero T and finite μ is important for the study of neutron stars.

Our approach for studying the problem:

Express the physical quantities we want to calculate in terms of various Green functions of QCD so that the problem reduces to the calculation of these Green functions. These Green functions can be calculated by various QCD methods/models (the Dyson-Schwinger equations, hard-thermal-loop/hard-dense-loop approximation, quasi-particle model, etc.)

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 6 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



★ A Formula for the EOS of QCD at Zero T and Finite μ

renormalized QCD partition function at zero T and finite μ :

$$\mathcal{Z}[\mu] = \int \mathcal{D}\bar{q}_R \mathcal{D}q_R \mathcal{D}A_R \exp \left\{ -S_R[\bar{q}_R, q_R, A_R] + \int d^4x \mu Z_2 \bar{q}_R(x) \gamma_4 q_R(x) \right\} \quad (1)$$

$S_R[\bar{q}_R, q_R, A_R]$: renormalized QCD action in Euclidean space

$Z_2 = Z_2(\zeta^2, \Lambda^2)$: quark wave-function renormalization constant

pressure density:

$$\mathcal{P}(\mu) = \frac{1}{V} \ln \mathcal{Z}[\mu] \quad (2)$$

quark number density:

$$\begin{aligned} \rho(\mu) &= \frac{\partial \mathcal{P}(\mu)}{\partial \mu} = \frac{1}{V} \frac{1}{\mathcal{Z}[\mu]} \frac{\partial \mathcal{Z}[\mu]}{\partial \mu} \\ &= \frac{1}{V} \frac{\int \mathcal{D}\bar{q}_R \mathcal{D}q_R \mathcal{D}A_R \int d^4x Z_2 \bar{q}_R(x) \gamma_4 q_R(x) \exp \{-S_R[\bar{q}_R, q_R, A_R; \mu]\}}{\int \mathcal{D}\bar{q}_R \mathcal{D}q_R \mathcal{D}A_R \exp \{-S_R[\bar{q}_R, q_R, A_R; \mu]\}} \end{aligned} \quad (3)$$

$$S_R[\bar{q}_R, q_R, A_R; \mu] \equiv S_R[\bar{q}_R, q_R, A_R] - \int d^4x \mu Z_2 \bar{q}_R(x) \gamma_4 q_R(x)$$

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 7 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



dressed quark propagator at finite μ

$$G_{Rij}[\mu](x, y) = \frac{\int \mathcal{D}\bar{q}_R \mathcal{D}q_R \mathcal{D}A_R q_{Ri}(x) \bar{q}_{Rj}(y) \exp \{-S_R[\bar{q}_R, q_R, A_R; \mu]\}}{\int \mathcal{D}\bar{q}_R \mathcal{D}q_R \mathcal{D}A_R \exp \{-S_R[\bar{q}_R, q_R, A_R; \mu]\}} \quad (4)$$

$$\text{Tr} \{G_R[\mu] \gamma_4\} = - \frac{\int \mathcal{D}\bar{q}_R \mathcal{D}q_R \mathcal{D}A_R \int d^4x \bar{q}_R(x) \gamma_4 q_R(x) \exp \{-S_R[\bar{q}_R, q_R, A_R; \mu]\}}{\int \mathcal{D}\bar{q}_R \mathcal{D}q_R \mathcal{D}A_R \exp \{-S_R[\bar{q}_R, q_R, A_R; \mu]\}} \quad (5)$$

Comparing (3) and (5) gives

$$\rho(\mu) = -\frac{Z_2}{\mathcal{V}} \text{Tr} \{G_R[\mu] \gamma_4\} = -N_c N_f Z_2 \int \frac{d^4p}{(2\pi)^4} \text{tr} \{G_R[\mu](p) \gamma_4\} \quad (6)$$

$\rho(\mu)$ is totally determined by the dressed quark propagator at finite μ

Integrating $\rho(\mu) = \frac{\partial \mathcal{P}(\mu)}{\partial \mu}$ gives

$$\mathcal{P}(\mu) = \mathcal{P}(\mu)|_{\mu=0} + \int_0^\mu d\mu' \rho(\mu') = \mathcal{P}(\mu)|_{\mu=0} - N_c N_f Z_2 \int_0^\mu d\mu' \int \frac{d^4p}{(2\pi)^4} \text{tr} \{G_R[\mu'](p) \gamma_4\} \quad (7)$$

Formula (7) is formally model-independent. When one actually applies it to calculate the EOS, one has to resort to various QCD methods/models.

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 8 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



★ Using Four Different Methods/Models to Calculate the EOS of QCD at Zero T and Finite μ

- Dyson-Schwinger Equation (DSE) of QCD
- Hard-Dense-Loop (HDL) Approximation
- Quasi-particle Model
- A Nonperturbative Approach Inspired by Chiral Perturbation Theory (χPT)

For details please refer to the following papers:

DSE:

H. S. Zong and W. M. Sun , *The calculation of the equation of state of QCD at finite chemical potential and zero temperature*, Phys. Rev. **D78**, 054001 (2008).

HDL:

Y. Jiang, H. Li, S. S. Huang, W. M. Sun and H. S. Zong *The equation of state and quark number susceptibility in hard-dense-loop approximation*, to appear in J. Phys. G.

Quasi-particle Model:

A. M. Zhao, J. Cao, L. J. Luo, W. M. Sun, and H. S. Zong, *The equation of state of quasi-particle model of quark-gluon plasma at finite chemical potential*, Mod. Phys. Lett. **A25**, 47-54 (2010).

A Nonperturbative Approach Inspired by χPT :

X. Y. Li, X. F. Lü, B. Wang, W. M. Sun, and H. S. Zong, *Study of the properties of cold dense nuclear matter based on a non-perturbative approach inspired by chiral perturbation theory*, Phys. Rev. **C80**, 034909 (2009).

[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 9 页 共 49 页

[返 回](#)


[全 屏 显 示](#)

[关 闭](#)


[退 出](#)



* DSE of QCD



DSE for quark propagator



DSE for quark-gluon vertex

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 10 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



$$\text{Diagrammatic representation of the gluon propagator DSE:}$$
$$G^{-1} = G_0^{-1} + \text{loop diagrams}$$

The equation shows the gluon propagator G^{-1} as the sum of its free part G_0^{-1} and various loop diagrams. The loops consist of gluons (wavy lines) and gluon-gluon vertices (black dots). There are six distinct loop configurations shown:

- A single loop with one gluon line entering and one exiting.
- A loop with two gluon lines entering and two exiting.
- A loop with three gluon lines entering and three exiting.
- A loop with four gluon lines entering and four exiting.
- A loop with five gluon lines entering and five exiting.
- A loop with six gluon lines entering and six exiting.

DSE for gluon propagator

[访问主页](#)

[标题页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 11 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



★ Calculation of EOS with Rainbow-Ladder DSE

rainbow-ladder quark DSE at zero μ

$$G^{-1}(p) = G_0^{-1}(p) + \frac{4}{3}g_s^2 \int \frac{d^4q}{(2\pi)^4} D_{\mu\nu}(p-q)\gamma_\mu G(q)\gamma_\nu$$

where $G_0^{-1}(p) = i\gamma \cdot p + m$ is the inverse of free quark propagator.

rainbow-ladder DSE at finite μ

$$G^{-1}[\mu](p) = (i\gamma \cdot p - \mu\gamma_4) + \frac{4}{3}g_s^2 \int \frac{d^4q}{(2\pi)^4} D_{\mu\nu}(p-q)\gamma_\mu G[\mu](q)\gamma_\nu \quad (8)$$

Adopting the approximation that the dressed gluon propagator is independent of μ and assuming the dressed quark propagator is analytic in the neighborhood of $\mu = 0$, one can prove

$$G_R^{-1}[\mu](p) = G_R^{-1}(\tilde{p}) = i\gamma \cdot \tilde{p}A(\tilde{p}^2) + B(\tilde{p}^2) \quad (9)$$

where $\tilde{p} \equiv (\vec{p}, p_4 + i\mu)$, $G_R^{-1}(p) = i\gamma \cdot pA(p^2) + B(p^2)$ is the inverse of the dressed quark propagator at zero μ .

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 12 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



For details of the proof, please see

H. S. Zong, L. Chang, F. Y. Hou, W. M. Sun, Y. X. Liu, Phys. Rev. C **71**, 015205 (2005)

F. Y. Hou, L. Chang, W. M. Sun, H. S. Zong, Y. X. Liu, Phys. Rev. C **72**, 034901 (2005)

H. T. Feng, F. Y. Hou, X. He, W. M. Sun, H. S. Zong, Phys. Rev. D **73**, 016004 (2006)

H. T. Feng, W. M. Sun, D. K. He, and H. S. Zong, Phys. Lett. B **661**, 57 (2008)

H. T. Feng, M. He, W. M. Sun, and H. S. Zong, Phys. Lett. B **688**, 178 (2010)

In the proof we have used the Taylor expansion of $G_R^{-1}[\mu](p)$ around $\mu = 0$. In the circle of convergence of its Taylor expansion (9) holds. But in fact (9) holds in the whole region in the complex μ plane where $G_R^{-1}[\mu](p)$ is analytic. This conclusion is based on a theorem in complex analysis:

Suppose two functions $f(z)$ and $g(z)$ are analytic in a common region D . If these two functions coincide in some portion $D' \subset D$, then they are equal everywhere in D .

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 13 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



For the dressed quark propagator at $\mu = 0$, we take the following model

(R. Alkofer et.al. Phys. Rev. D **70**, 014014 (2004).)

$$G_R(p) = Z_2^{-1}(\zeta^2, \Lambda^2) \sum_{j=1}^{n_P} \left\{ \frac{r_j}{i \not{p} + m_j} + \frac{r_j}{i \not{p} + m_j^*} \right\} \quad (10)$$

where $m_j = a_j + ib_j$ are complex mass scales, r_j are real coefficients. The renormalization is chosen at $\zeta^2 = 16 \text{ GeV}^2$.

$$\begin{aligned} G_R[\mu](p) &= Z_2^{-1}(\zeta^2, \Lambda^2) \sum_{j=1}^{n_P} \left\{ \frac{r_j}{i \not{p} + m_j} + \frac{r_j}{i \not{p} + m_j^*} \right\} \\ &= Z_2^{-1}(\zeta^2, \Lambda^2) \sum_{j=1}^{n_P} \left\{ \frac{r_j(-i \not{p} + m_j)}{\not{p}^2 + m_j^2} + \frac{r_j(-i \not{p} + m_j^*)}{\not{p}^2 + m_j^{*2}} \right\} \end{aligned} \quad (11)$$

From this we obtain the quark number density

$$\begin{aligned} \rho(\mu) &= 4iN_c N_f \int \frac{d^4 p}{(2\pi)^4} \sum_{j=1}^{n_P} \left\{ \frac{r_j(p_4 + i\mu)}{\not{p}^2 + (p_4 + i\mu)^2 + m_j^2} + \frac{r_j(p_4 + i\mu)}{\not{p}^2 + (p_4 + i\mu)^2 + m_j^{*2}} \right\} \\ &= \frac{4iN_c N_f}{(2\pi)^4} \sum_{j=1}^{n_P} \left\{ r_j \int d^4 p \frac{p_4 + i\mu}{\not{p}^2 + (p_4 + i\mu)^2 + m_j^2} + r_j \int d^4 p \frac{p_4 + i\mu}{\not{p}^2 + (p_4 + i\mu)^2 + m_j^{*2}} \right\} \end{aligned} \quad (12)$$

[访问主页](#)

[标 题 页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 14 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



the calculation of the integral in (12)

$$\int d^4 p \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m_j^2} = \int d\vec{p} \int_{-\infty}^{+\infty} dp_4 \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m_j^2}$$

the integral of p_4 can be written as

$$\int_{-\infty}^{+\infty} dp_4 \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m_j^2} = \int_{-\infty+i\mu}^{+\infty+i\mu} dz \frac{z}{z^2 + \vec{p}^2 + m_j^2} \quad (13)$$

(13) can be calculated by method of contour integral

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 15 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)




choose the following rectangular contour

the pole of the function $f(z) = \frac{z}{z^2 + \vec{p}^2 + m_j^2}$ in the upper complex z plane:

$$\begin{aligned} z_j &= \operatorname{sgn}(-\beta_j) \sqrt{\frac{\sqrt{(\vec{p}^2 + \alpha_j)^2 + \beta_j^2} - \vec{p}^2 - \alpha_j}{2}} + i \sqrt{\frac{\sqrt{(\vec{p}^2 + \alpha_j)^2 + \beta_j^2} + \vec{p}^2 + \alpha_j}{2}} \\ &= \chi_j(\vec{p}) + i\omega_j(\vec{p}) \end{aligned}$$

where $m_j^2 \equiv \alpha_j + \beta_j i$



- (i) $\mu < \omega_j(\vec{p})$ The contour contains no poles, the integral along the contour vanishes. $f(z)$ is an odd function, the integral along $[-R, R]$ vanishes. So one has

$$\int_{-\infty+i\mu}^{+\infty+i\mu} dz f(z) = 0, \quad \mu < \omega_j(\vec{p})$$

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)


第 16 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



(ii) $\mu > \omega_j(\vec{p})$ The contour contains one pole. According to residue theorem, one has

$$-\int_{-\infty+i\mu}^{+\infty+i\mu} dz f(z) = 2\pi i \operatorname{Res}(f(z), z = z_j) = \pi i, \quad \mu > \omega_j(\vec{p})$$

(i) , (ii) can be combined to be written as

$$\int_{-\infty}^{+\infty} dp_4 \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m_j^2} = \int_{-\infty+i\mu}^{+\infty+i\mu} dz f(z) = -\pi i \theta(\mu - \omega_j(\vec{p})) \quad (14)$$

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 17 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



when $\mu < \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}$, one has $\omega_j(\vec{p}) > \mu$ irrespective the value of \vec{p} ;

when $\mu \geq \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}$, $\omega_j(\vec{p}) < \mu$ if and only if $|\vec{p}| < (\mu^2 - \frac{\beta_j^2}{4\mu^2} - \alpha_j)^{1/2}$. So

$$\begin{aligned} \int d^4p \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m_j^2} &= -\pi i \int d\vec{p} \theta(\mu - \omega_j(\vec{p})) \\ &= \begin{cases} 0, & \text{when } \mu < \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}} \\ -\frac{4\pi^2 i}{3} (\mu^2 - \frac{\beta_j^2}{4\mu^2} - \alpha_j)^{3/2}, & \text{when } \mu \geq \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}} \end{cases} \\ &= -\frac{4\pi^2 i}{3} \theta(\mu - \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}) (\mu^2 - \frac{\beta_j^2}{4\mu^2} - \alpha_j)^{3/2} \end{aligned} \quad (15)$$

Make the replacement $\beta_j \rightarrow -\beta_j$ in (15), one obtains

$$\int d^4p \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m_j^{*2}} = -\frac{4\pi^2 i}{3} \theta(\mu - \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}) (\mu^2 - \frac{\beta_j^2}{4\mu^2} - \alpha_j)^{3/2} \quad (16)$$

Substituting (15),(16) into (12) gives

$$\rho(\mu) = \frac{2N_c N_f}{3\pi^2} \sum_{j=1}^{n_p} r_j \theta(\mu - \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}) (\mu^2 - \frac{\beta_j^2}{4\mu^2} - \alpha_j)^{3/2} \quad (17)$$

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 18 页 共 49 页

[返 回](#)

[全屏显示](#)


[关 闭](#)

[退 出](#)

In numerical calculations, one employs three sets of parameters (taken from R. Alkofer et.al. Phys. Rev. D **70**, 014014 (2004)), representing three forms of the quark propagator: three real poles(3R), two pairs of complex conjugate poles(2CC), one real pole and a pair of complex conjugate poles(1R1CC)

Parametrization	r_1	a_1 (GeV)	b_1 (GeV)	r_2	a_2 (GeV)	b_2 (GeV)	r_3	a_3 (GeV)
2CC	0.360	0.351	0.08	0.140	-0.899	0.463
1R1CC	0.354	0.377	...	0.146	-0.91	0.45
3R	0.365	0.341	...	1.2	-1.31	...	-1.06	-1.40

the dependence of the quark number density on μ



when μ is less than a critical value μ_0 (for 2CC, 1R1CC and 3R parametrizations, μ_0 equals 351 MeV, 377 MeV and 341 MeV, respectively), the quark number density vanishes identically. $\mu = \mu_0$ is a singularity.



[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 19 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



It is a model-independent result that some singularity exists at $\mu = \mu_0$ and $T = 0$
(see M. A. Halasz et. al. Phys. Rev. D **58**, 096007 (1998))

Argument: The QCD partition function is

$$Z = \sum_{\alpha} \exp\left\{-\frac{E_{\alpha} - \mu N_{\alpha}}{T}\right\}$$

In the limit $T \rightarrow 0$, the state with lowest $E_{\alpha} - \mu N_{\alpha}$ gives an exponentially dominant contribution to the partition function.

When $\mu = 0$, this is the state with $N = 0$ and $E = 0$, i.e., the vacuum.

Introducing

$$\mu_0 \equiv \min_{\alpha} \frac{E_{\alpha}}{|N_{\alpha}|}$$

When $\mu < \mu_0$, the state with the lowest $E_{\alpha} - \mu N_{\alpha}$ is still the vacuum. Therefore, at $T = 0$

$$\rho(\mu) = 0, \quad \mu < \mu_0$$

For pure QCD (electromagnetic interaction being switched off), μ_0 is estimated to be $\frac{m_N - 16 \text{ MeV}}{N_c} = 307 \text{ MeV}$ (for details, please see the above reference).

[访问主页](#)

[标题页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 20 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)

From (17) one can obtain



$$\begin{aligned}
 \int_0^\mu d\mu' \rho(\mu') &= \frac{2N_c N_f}{3\pi^2} \sum_{j=1}^{n_p} r_j \int_0^\mu d\mu' \theta\left(\mu' - \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}\right) (\mu'^2 - \frac{\beta_j^2}{4\mu'^2} - \alpha_j)^{3/2} \\
 &= \frac{2N_c N_f}{3\pi^2} \sum_{j=1}^{n_p} r_j \theta\left(\mu - \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}\right) \int_{\sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}}^\mu d\mu' (\mu'^2 - \frac{\beta_j^2}{4\mu'^2} - \alpha_j)^{3/2} \\
 &= \frac{2N_c N_f}{3\pi^2} \sum_{j=1}^{n_p} r_j \theta\left(\mu - \sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}\right) I(\mu; \alpha_j, \beta_j).
 \end{aligned} \tag{18}$$

where $I(\mu; \alpha_j, \beta_j)$ is

$$\begin{aligned}
 I(\mu; \alpha_j, \beta_j) &\equiv \int_{\sqrt{\frac{\alpha_j + \sqrt{\alpha_j^2 + \beta_j^2}}{2}}}^\mu d\mu' \left(\mu'^2 - \frac{\beta_j^2}{4\mu'^2} - \alpha_j\right)^{3/2} \\
 &= \frac{3(\alpha_j^2 - \beta_j^2)}{16} \ln \frac{\sqrt{\mu^2 - \alpha_j/2 + \sqrt{\alpha_j^2 + \beta_j^2}/2} + \sqrt{\mu^2 - \alpha_j/2 - \sqrt{\alpha_j^2 + \beta_j^2}/2}}{\sqrt{\mu^2 - \alpha_j/2 + \sqrt{\alpha_j^2 + \beta_j^2}/2} - \sqrt{\mu^2 - \alpha_j/2 - \sqrt{\alpha_j^2 + \beta_j^2}/2}} \\
 &\quad + \frac{3\alpha_j |\beta_j|}{4} \arctan \frac{\sqrt{(\sqrt{\alpha_j^2 + \beta_j^2} - \alpha_j)(\mu^2 - \sqrt{\alpha_j^2 + \beta_j^2}/2 - \alpha_j/2)}}{\sqrt{(\sqrt{\alpha_j^2 + \beta_j^2} + \alpha_j)(\mu^2 + \sqrt{\alpha_j^2 + \beta_j^2}/2 - \alpha_j/2)}} \\
 &\quad + \frac{\mu^2}{4} \sqrt{\mu^4 - \alpha_j \mu^2 - \beta_j^2/4} - \frac{5\alpha_j}{8} \sqrt{\mu^4 - \alpha_j \mu^2 - \beta_j^2/4} + \frac{\beta_j^2}{8} \frac{\sqrt{\mu^4 - \alpha_j \mu^2 - \beta_j^2/4}}{\mu^2}.
 \end{aligned} \tag{19}$$

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 21 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)

(19)



$\mathcal{P}(\mu)|_{\mu=0}$ can be calculated from thw CJT effective action.

$$\begin{aligned}\mathcal{P}(\mu)|_{\mu=0} &= 2N_c N_f \int \frac{d^4 p}{(2\pi)^4} \left\{ \ln \left[\frac{A^2(p^2)p^2 + B^2(p^2)}{p^2} \right] - \frac{p^2 A(p^2)[A(p^2) - 1] + B^2(p^2)}{p^2 A^2(p^2) + B^2(p^2)} \right\} \\ &= -2N_c N_f \int \frac{d^4 p}{(2\pi)^4} \left\{ \ln \left[p^2 \left(p^2 \sigma_v^2(p^2) + \sigma_s^2(p^2) \right) \right] + 1 + p^2 \sigma_v(p^2) \right\} \quad (20)\end{aligned}$$

其中 $G(p) = \frac{1}{i\gamma \cdot p A(p^2) + B(p^2)} \equiv i\gamma \cdot p \sigma_v(p^2) + \sigma_s(p^2)$ is the unrenormalized dressed quark propagator at $\mu = 0$.

For the model quark propagator we use

$$\sigma_v(p^2) = - \sum_{j=1}^{n_P} \left(\frac{r_j}{p^2 + m_j^2} + \frac{r_j}{p^2 + m_j^{*2}} \right), \quad \sigma_s(p^2) = \sum_{j=1}^{n_P} \left(\frac{r_j m_j}{p^2 + m_j^2} + \frac{r_j m_j^*}{p^2 + m_j^{*2}} \right) \quad (21)$$

From (20), (21) and the model parameters given before, one can calculate $\mathcal{P}(\mu)|_{\mu=0}$.

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 22 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



Comparison of our EOS with the cold, perturbative EOS of QCD

In E. S. Fraga, R. D. Pisarski, J. Schaffner-Bielich, Phys. Rev. **D 63**, 121702(R) (2001), the following EOS was obtained:

$$\mathcal{P}_{FPS}(\mu) = \frac{N_f \mu^4}{4\pi^2} \left\{ 1 - 2 \left(\frac{\alpha_s}{\pi} \right) - \left[G + N_f \ln \frac{\alpha_s}{\pi} + \left(11 - \frac{2}{3} N_f \right) \ln \frac{\bar{\Lambda}}{\mu} \right] \left(\frac{\alpha_s}{\pi} \right)^2 \right\} \quad (22)$$

Here we adopt \overline{MS} scheme, $G = G_0 - 0.536 N_f + N_f \ln N_f$, $G_0 = 10.374 \pm 0.13$, $\bar{\Lambda}$ is the renormalization subtraction point.

$$\alpha_s(\bar{\Lambda}) = \frac{4\pi}{\beta_0 u} \left[1 - \frac{2\beta_1 \ln(u)}{\beta_0^2} + \frac{4\beta_1^2}{\beta_0^4 u^2} \left(\left(\ln(u) - \frac{1}{2} \right)^2 + \frac{\beta_2 \beta_0}{8\beta_1^2} - \frac{5}{4} \right) \right]$$

where $u = \ln(\bar{\Lambda}^2/\Lambda_{\overline{MS}}^2)$, $\beta_0 = 11 - 2N_f/3$, $\beta_1 = 51 - 19N_f/3$, $\beta_2 = 2857 - 5033N_f/9 + 325N_f^2/27$.

For $N_f = 3$, $\Lambda_{\overline{MS}} = 365$ MeV. The ratio $\bar{\Lambda}/\mu$ is taken to be 2.

EOS (22) is only valid in the chirally symmetric phase ($\mu > \mu_\chi$)

[访问主页](#)

[标题页](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

第 23 页 共 49 页

[返 回](#)


[全屏显示](#)

[关 闭](#)

[退 出](#)



Comparison of our EOS with the perturbative EOS (when making the comparison we have neglected the constant term $\mathcal{P}(\mu)|_{\mu=0}$)



[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 24 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



★ Calculation of EOS with HDL Approximation

At sufficiently high chemical potential, the HDL approximation is a good approximation to QCD.
quark propagator in HDL approximation

$$G_R(p) = -Z_2^{-1} \frac{1}{D_+(p)} \frac{\gamma_4 + i\hat{\mathbf{p}} \cdot \vec{\gamma}}{2} - Z_2^{-1} \frac{1}{D_-(p)} \frac{\gamma_4 - i\hat{\mathbf{p}} \cdot \vec{\gamma}}{2}, \quad (23)$$

where $\hat{\mathbf{p}} = \vec{p}/|\vec{p}|$, $p_4 = (2n+1)\pi T$ ($n \in \mathbf{Z}$) are the fermion Matsubara frequencies

$$D_{\pm}(p) = -ip_4 \pm |\vec{p}| + \mu + \frac{m_q^2}{|\vec{p}|} \left[Q_0 \left(\frac{ip_4 - \mu}{|\vec{p}|} \right) \mp Q_1 \left(\frac{ip_4 - \mu}{|\vec{p}|} \right) \right] \quad (24)$$

$m_q \equiv g\sqrt{(T^2 + \mu^2/\pi^2)/6}$ is the quark thermal mass (g is the strong coupling constant)

[访问主页](#)

[标题页](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

第 25 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)




From this one obtains the baryon number density under HDL approximation

$$\rho(\mu, T) = 2N_c N_f \int \frac{d^3 \vec{p}}{(2\pi)^3} T \sum_n \left[\frac{1}{D_+} + \frac{1}{D_-} \right] \quad (25)$$

The summation over Matsubara frequencies can be calculated by contour integral method

$$\begin{aligned} & T \sum_n \left[\frac{1}{D_+} + \frac{1}{D_-} \right] + \int_{-i|\vec{p}| - i\mu}^{i|\vec{p}| - i\mu} \frac{dp_4}{2\pi} \text{Disc} \left[\frac{1}{D_+} + \frac{1}{D_-} \right] \frac{1}{2} \tanh \frac{ip_4}{2T} \\ &= \int_{C_1 \cup C_2} \frac{dp_4}{2\pi} \left[\frac{1}{D_+} + \frac{1}{D_-} \right] \frac{1}{2} \tanh \frac{ip_4}{2T}. \end{aligned} \quad (26)$$

the integral contour



[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 26 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



The calculation of the integral:

Define

$$f(p_4) := \left[\frac{1}{D_+} + \frac{1}{D_-} \right] \frac{1}{2} \tanh \frac{ip_4}{2T}$$

and closing C_1 and C_2 with large half circles, one obtains the following

$$\int_{C_1 \cup C_2} \frac{dp_4}{2\pi} \left[\frac{1}{D_+} + \frac{1}{D_-} \right] \frac{1}{2} \tanh \frac{ip_4}{2T} = \frac{1}{2\pi} (-2\pi i) \sum_j \text{Res}\{f(\zeta_j)\}, \quad (27)$$

where ζ_j are the poles of $f(p_4)$ located in the imaginary axis. The poles of $1/D_+$ are

$$\zeta_1 = -i(\omega_+ + \mu), \quad \text{Res} \left\{ \frac{1}{D_+(\zeta_1)} \right\} = iZ_+ \quad (28)$$

$$\zeta_2 = i(\omega_- - \mu), \quad \text{Res} \left\{ \frac{1}{D_+(\zeta_2)} \right\} = iZ_- \quad (29)$$

and the poles of $1/D_-$ are

$$\zeta_3 = i(\omega_+ - \mu), \quad \text{Res} \left\{ \frac{1}{D_-(\zeta_3)} \right\} = iZ_+ \quad (30)$$

$$\zeta_4 = -i(\omega_- + \mu), \quad \text{Res} \left\{ \frac{1}{D_-(\zeta_4)} \right\} = iZ_-, \quad (31)$$

[访问主页](#)

[标题页](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

第 27 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



$\omega_{\pm}(|\vec{p}|)(>|\vec{p}|)$ are the solutions of the equations

$$\frac{|\vec{p}|(\omega_{+} - |\vec{p}|)}{m_q^2} - 1 = \frac{1}{2} \left(1 - \frac{\omega_{+}}{|\vec{p}|}\right) \ln \frac{\omega_{+} + |\vec{p}|}{\omega_{+} - |\vec{p}|}, \quad (32)$$

$$\frac{|\vec{p}|(\omega_{-} + |\vec{p}|)}{m_q^2} + 1 = \frac{1}{2} \left(1 + \frac{\omega_{-}}{|\vec{p}|}\right) \ln \frac{\omega_{-} + |\vec{p}|}{\omega_{-} - |\vec{p}|} \quad (33)$$

and $Z_{\pm}(|\vec{p}|)$ are

$$Z_{\pm}(|\vec{p}|) = \frac{\omega_{\pm} - |\vec{p}|^2}{2m_q^2}. \quad (34)$$

From these results one can calculate the quasi-particle contribution:

$$\begin{aligned} & \int_{C_1 \cup C_2} \frac{dp_4}{2\pi} \left[\frac{1}{D_{+}} + \frac{1}{D_{-}} \right] \frac{1}{2} \tanh \frac{ip_4}{2T} = \frac{1}{2\pi} (-2\pi i) \sum_j \text{Res}\{f(\zeta_j)\} \\ &= -\frac{i}{2} \left[iZ_{+} \tanh \frac{\omega_{+} + \mu}{2T} + iZ_{-} \tanh \frac{-(\omega_{-} - \mu)}{2T} + iZ_{+} \tanh \frac{-(\omega_{+} - \mu)}{2T} + iZ_{-} \tanh \frac{\omega_{-} + \mu}{2T} \right]. \end{aligned} \quad (35)$$

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 28 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



The cut contribution can be expressed as

$$\int_{-i|\vec{p}| - i\mu}^{i|\vec{p}| - i\mu} \frac{dp_4}{2\pi} \text{Disc} \left(\frac{1}{D_+} + \frac{1}{D_-} \right) \frac{1}{2} \tanh \frac{ip_4}{2T} = \int_{-|\vec{p}|}^{|\vec{p}|} \frac{d\omega}{2\pi} [\rho_+(\omega) + \rho_-(\omega)] \frac{1}{2} \tanh \frac{\omega - \mu}{2T},$$

where $\rho_{\pm}(\omega, |\vec{p}|)$ are the familiar spectral functions of the quark propagator ($x \equiv \omega/|\vec{p}|$):

$$\begin{aligned} \rho_{\pm}(\omega, |\vec{p}|) &= 2\pi [Z_{\pm}(|\vec{p}|) \delta(\omega - \omega_{\pm}(|\vec{p}|)) + Z_{\mp}(|\vec{p}|) \delta(\omega + \omega_{\mp}(|\vec{p}|))] \\ &\quad + \frac{\pi m_q^2 (1 \mp x) \theta(1 - x^2)}{|\vec{p}|^3} \left\{ \left[1 \mp x + \frac{m_q^2}{|\vec{p}|^2} \pm \frac{m_q^2}{2|\vec{p}|^2} (1 \mp x) \ln \frac{1+x}{1-x} \right]^2 \right. \\ &\quad \left. + \frac{\pi^2 m_q^4}{4|\vec{p}|^4} (1 \mp x)^2 \right\}^{-1}. \end{aligned} \quad (36)$$

Combining the quasi-particle and the Landau damping contributions one can obtain

$$\begin{aligned} T \sum_n \left[\frac{1}{D_+} + \frac{1}{D_-} \right] &= Z_+ \left[n_F(\omega_+ - \mu) - n_F(\omega_+ + \mu) \right] + Z_- \left[n_F(\omega_- + \mu) - n_F(\omega_- - \mu) \right] \\ &\quad - \frac{1}{2} \int_{-|\vec{p}|}^{|\vec{p}|} \frac{d\omega}{2\pi} [\rho_+(\omega) + \rho_-(\omega)] [1 - 2n_F(\omega - \mu)], \end{aligned} \quad (37)$$

where $n_F(\omega)$ is the Fermi distribution function

$$n_F(\omega) = \frac{1}{\exp(\omega/T) + 1}. \quad (38)$$

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 29 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



Therefore the quark number density at finite μ and T is

$$\begin{aligned}\rho(\mu, T) = & 2N_c N_f \int \frac{d^3 \vec{p}}{(2\pi)^3} \left\{ Z_+ \left[n_F(\omega_+ - \mu) - n_F(\omega_+ + \mu) \right] \right. \\ & + Z_- \left[n_F(\omega_- + \mu) - n_F(\omega_- - \mu) \right] \left. \right\} \\ & - N_c N_f \int \frac{d^3 \vec{p}}{(2\pi)^3} \int_{-\vec{p}}^{|\vec{p}|} \frac{d\omega}{2\pi} [\rho_+(\omega) + \rho_-(\omega)] [1 - 2n_F(\omega - \mu)].\end{aligned}\quad (39)$$

In the limit of $T \rightarrow \infty$, ρ tends to the free quark gas result ($Z_+ \rightarrow 1$, $Z_- \rightarrow 0$ and the integral is dominated by $p \sim \omega \sim T$ region)

[访问主页](#)

[标 题 页](#)

[◀](#) | [▶](#)

[◀](#) | [▶](#)

第 30 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)


[退 出](#)

Baryon number density at zero T and finite μ :

$$\begin{aligned}\rho(\mu) = & 2N_c N_f \int \frac{d^3 \vec{p}}{(2\pi)^3} [\theta(\mu - \omega_+) Z_+ - \theta(\mu - \omega_-) Z_-] \\ & + N_c N_f \int \frac{d^3 \vec{p}}{(2\pi)^3} \int_{-\|\vec{p}\|}^{\|\vec{p}\|} \frac{d\omega}{2\pi} [\rho_+(\omega) + \rho_-(\omega)] [1 - 2\theta(\omega - \mu)].\end{aligned}\quad (40)$$

EOS under HDL approximation (here the constant term $\mathcal{P}(\mu)|_{\mu=0}$ is neglected)

$$\begin{aligned}\mathcal{P}(\mu) = & 2N_c N_f \int_0^\mu d\mu' \int \frac{d^3 \vec{p}}{(2\pi)^3} [\theta(\mu' - \omega_+) Z_+ - \theta(\mu' - \omega_-) Z_-] \\ & + N_c N_f \int_0^\mu d\mu' \int \frac{d^3 \vec{p}}{(2\pi)^3} \int_{-\|\vec{p}\|}^{\|\vec{p}\|} \frac{d\omega}{2\pi} [\rho_+(\omega) + \rho_-(\omega)] [1 - 2\theta(\omega - \mu')].\end{aligned}\quad (41)$$



访问主页

标题页

◀ ▶

◀ ▶

第 31 页 共 49 页

返 回


全屏显示

关 闭

退 出



comparison of HDL EOS with the perturbative EOS



[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 32 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



★ Calculation of EOS with the Quasi-particle Model

• Introduction to quasi-particle model

In quasi-particle model, the interacting particle system is effectively regarded as an ideal gas of non-interacting quasi-particles with temperature and density dependent effective mass.

In quasi-particle model, the quark propagator has the form of a free fermion propagator with the effective mass

$$G^{-1}[\mu](\vec{p}, \omega_n) = i\vec{\gamma} \cdot \vec{p} + i\gamma_4(\omega_n + i\mu) + m(\mu, T)$$

$\omega_n = (2n + 1)\pi T$, $n \in \mathbf{Z}$ are the fermion Matsubara frequencies.

The effective mass is taken to be (X.P. Zhang, M. Kang, X.W. Liu, and S.H. Yang, Phys. Rev. C **72**, 025809 (2005)):

$$m^2(T, \mu) = aT^2 + \frac{N_c^2 - 1}{8N_c}(T^2 + \frac{\mu^2}{\pi^2})g^2(T, \mu) \quad (42)$$

$g^2(T, \mu)$ is the effective strong coupling constant.

At $T = 0$, the effective mass and effective strong coupling constant are

$$m^2(\mu) = \frac{\mu^2}{3\pi^2}g^2(0, \mu), \quad g^2(0, \mu) = \frac{16\pi^2}{9 \ln(\frac{\mu+T_s}{T_c\pi\sqrt{\lambda}})^2} \quad (43)$$

where the parameters are $\lambda = 6.6$, $T_c = 170$ MeV, $T_s = -0.78T_c$

In the limit $T \rightarrow 0$, discrete Matsubara frequency $\omega_n \Rightarrow$ continuous variable p_4

访问主页

标题页

◀ ▶

◀ ▶

第 33 页 共 49 页

返 回

全屏显示

关 闭

退 出

baryon number density at zero T and finite μ

$$\rho(\mu) = 4iN_c N_f \int \frac{d^4 p}{(2\pi)^4} \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m^2(\mu)} \quad (44)$$

the calculation of the integral

$$\int d^4 p \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m(\mu)^2} = \int d\vec{p} \int_{-\infty}^{+\infty} dp_4 \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m(\mu)^2}$$

The integral of p_4 can be calculated by the contour integral method. The result is

$$\int_{-\infty}^{+\infty} dp_4 \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m(\mu)^2} = -\pi i \theta(\mu - \omega_j(\vec{p}, \mu)), \quad \omega_j(\vec{p}, \mu) = \sqrt{\vec{p}^2 + m(\mu)^2}$$

Using

$$\theta(\mu - \omega_j(\vec{p}, \mu)) = \begin{cases} 1, & \text{when } |\vec{p}| < (\mu^2 - m(\mu)^2)^{1/2} \\ 0, & \text{when } |\vec{p}| > (\mu^2 - m(\mu)^2)^{1/2}. \end{cases}$$

one obtains

$$\begin{aligned} \int d^4 p \frac{p_4 + i\mu}{\vec{p}^2 + (p_4 + i\mu)^2 + m(\mu)^2} &= -\pi i \int d\vec{p} \theta(\mu - \omega_j(\vec{p}, \mu)) \\ &= \begin{cases} 0, & \text{when } \mu < m(\mu) \\ -\frac{4\pi^2 i}{3} (\mu^2 - m(\mu)^2)^{3/2}, & \text{when } \mu \geq m(\mu) \end{cases} \\ &= -\frac{4\pi^2 i}{3} \theta(\mu - m(\mu)) (\mu^2 - m(\mu)^2)^{3/2} \end{aligned}$$



访问主页

标 题 页

◀ ▶

◀ ▶

第 34 页 共 49 页

返 回

全 屏 显 示


关 闭

退 出



Thus one has

$$\rho(\mu) = \frac{3}{\pi^2} \theta(\mu - m(\mu)) (\mu^2 - m(\mu)^2)^{3/2} \quad (45)$$



The obtained baryon number density differs significantly with the free fermion gas result. This difference comes from the chemical potential dependence of the effective mass.

critical chemical potential: $\mu_0 = 241$ MeV, This is comparable to the result $\mu_0 = 307$ MeV in M. A. Halasz et. al. Phys. Rev. D **58**, 096007 (1998).

[访问主页](#)

[标题页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 35 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)


[退 出](#)



EOS:

$$\mathcal{P}(\mu) = \mathcal{P}(\mu)|_{\mu=0} + \frac{3}{\pi^2} \int_0^\mu d\mu' \theta(\mu' - m(\mu')) (\mu'^2 - m(\mu')^2)^{3/2} \quad (46)$$

Comparison of the EOS in the quasi-particle model with the perturbative EOS of QCD (the constant term $\mathcal{P}(\mu)|_{\mu=0}$ is neglected)



The form of the EOS in the quasi-particle model depends on the form of the effective mass in the quasi-particle model and the parameters therein. The parameters in the effective mass employed in our calculation has some arbitrariness. It is expected that the result of EOS can give some constraints on the choice of these parameters.

[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 36 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



★ Calculation of EOS with a Nonperturbative Approach inspired by χPT

χPT is an efficient method of studying the evolution properties of cold, dense strongly interacting matter with the chemical potential in low energy regime (where chiral symmetry is spontaneously broken)

Limitation of χPT : it is only valid in the low energy regime. J.A.Oller and E.Oset developed a nonperturbative method, extending the idea of χPT beyond its range of validity.
(J.A.Oller and E.Oset, Nucl. Phys. A 620, 438 (1997))

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 37 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)

- Effective chiral Lagrangian at zero T and finite baryon chemical potential

baryon number density operator: $\frac{1}{3}q^\dagger q$

baryon number density: $\langle \frac{1}{3}q^\dagger q \rangle$

The hadron system can be described by an effective Lagrangian containing all possible terms consistent with the symmetry principles. One can calculate the vacuum expectation value of the quark operators from χPT .

The calculation of $\langle \bar{q}\gamma_\mu q \rangle, \langle \bar{q}\gamma_5\gamma_\mu q \rangle, \langle \bar{q}q \rangle, \langle \bar{q}\gamma_5 q \rangle$ in χPT :

Introducing external fields: a_μ, v_μ, s, p

$$\mathcal{L}_{ext}^{QCD} = \mathcal{L}_0^{QCD} + \bar{q}\gamma^\mu(v_\mu + \gamma_5 a_\mu)q - \bar{q}\gamma^\mu(s - i\gamma_5 p)q$$

In low energy regime the effective Lagrangian contains all possible terms consistent with symmetry principles. The generating functional derived from this Lagrangian is equivalent to the generating functional derived from QCD. The vacuum expectation value of the quark current can be obtained by differentiation of the generating functional with respect to the external fields.

The calculation of baryon number density $\langle \frac{1}{3}q^\dagger q \rangle$

Introduction of baryon chemical potential:

Introduce the external field b_μ and do the replacement $\partial_\mu q \rightarrow \partial_\mu q - B^q b_\mu q = \partial_\mu q - \frac{i}{3}b_\mu q$

In the last step of calculation we set $b_\mu = \delta_{\mu 0}\mu_b$, where μ_b is the baryon chemical potential.

Because of $SU(2)_f$ symmetry, u, d quark have equal mass and chemical potential.



[访问主页](#)

[标 题 页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 38 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)

baryon number density can be obtained by differentiation of the generating functional:

$$\langle \frac{1}{3}q^+q \rangle = \frac{\partial Z[b_\mu]}{\partial b_0}|_{b_0=\mu_b, b_i=0} = \frac{\partial Z[\mu_b]}{\partial \mu_b}$$

introduction of chemical potential at the nucleon level: in $\mathcal{L}_{\pi N}$ replace $\partial_\mu \psi$ by

$$\partial_\mu \psi \rightarrow \partial_\mu \psi - iB^N b_\mu \psi = \partial_\mu \psi - i\delta_{0\mu} \mu_b \psi$$

Effective Lagrangian: for pionic part one only needs to consider the leading order term:

$$\mathcal{L}_\pi^{(2)} = \frac{F_\pi^2}{4} Tr(\partial_\mu U \partial^\mu U^+) + \frac{F_\pi^2}{4} Tr(\chi U^+ + U \chi^+)$$

$$U = \exp\left(\frac{i\vec{\pi} \cdot \vec{\tau}}{F_\pi}\right), \quad \chi = 2B_0(s + ip)$$

in our calculation $p = 0$, s = mass matrix of u, d quark

under $SU(2)_f$, $\chi = 2B_0 M_q = 2B_0 \hat{m} = M_\pi^2$

for πN part, to $O(p^4)$ order

$$\mathcal{L}_{\pi N}^{(1)} = \bar{\psi}(i \not{D} - m)\psi + \frac{1}{2}g_A \bar{\psi} \not{\mu} \gamma_5 \psi,$$

$$\mathcal{L}_{\pi N}^{(2)} = c_1 \langle \chi_+ \rangle \bar{\psi} \psi - \frac{c_2}{4m^2} \langle u_\mu u_\nu \rangle (\bar{\psi} D^\mu D^\nu \psi + h.c.) + \frac{c_3}{2} u_\mu u^\mu \bar{\psi} \psi + \dots,$$

$$\mathcal{L}_{\pi N}^{(4)} = -\frac{e_1}{16} \langle \chi_+ \rangle^2 \bar{\psi} \psi + \dots$$

where

$$D_\mu \psi = \partial_\mu \psi - i\delta_{0\mu} \mu_b \psi + \Gamma_\mu \psi,$$

$$\Gamma_\mu = \frac{1}{2}[u^+, \partial_\mu u],$$

$$u^2 = U, u_\mu = iu^+ \partial_\mu U u^+$$



访问主页

标题页

◀ ▶

◀ ▶

第 39 页 共 49 页

返 回

全屏显示

关闭

退出



The numerical values of m , M_π , F_π , g_A , c_1, c_2, c_3, e_1 used in our calculation. The values of c_1, c_2, c_3 are the same as those in V. Bernard, Prog. Part. Nucl. Phys. **60**, 82, 2007. The values of m and e_1 are obtained through fitting the nucleon mass.

$M_\pi [MeV]$	$m [MeV]$	$F_\pi [MeV]$	g_A	$c_1 [GeV^{-3}]$	$c_2 [GeV^{-3}]$	$c_3 [GeV^{-3}]$	$e_1 [GeV^{-2}]$
137	896	92.4	1.27	-0.90×10^{-3}	3.3×10^{-3}	-4.7×10^{-3}	1×10^{-9}

[访问主页](#)

[标 题 页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 40 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



- The calculation of baryon number density and EOS

$$\mathcal{L}_{\pi N} = \bar{\psi} \hat{K} \psi, \quad \hat{K} = i\gamma_\mu \partial_\mu + \gamma_0 \mu_b - m + \hat{O}$$

baryon number density

$$\begin{aligned} n(\mu_b) &= \langle \frac{1}{3} q^+ q \rangle = \frac{\partial}{\partial \mu_b} \int D\bar{\psi} D\psi DU e^{i \int d^4x \mathcal{L}_{eff}} \\ &= \frac{\partial}{\partial \mu_b} \int DU (Det K) e^{i \int d^4x \mathcal{L}_\pi} \\ &= \int DUTr(K^{-1} \frac{\partial}{\partial \mu_b} K) e^{i \int d^4x \mathcal{L}_\pi} \end{aligned}$$

where $\mathcal{L}_{eff} = \mathcal{L}_{\pi N} + \mathcal{L}_\pi$.

K, O are the results of the path integral after integrating out the nucleon field.

The most important contribution to baryon number density is from $Tr(K^{-1} \gamma_0)$, the contribution of $Tr(K^{-1} \frac{\partial}{\partial \mu_b} O)$ can be neglected.

$$\begin{aligned} n(\mu_b) &\doteq \int DUTr(K^{-1} \gamma_0) e^{i \int d^4x \mathcal{L}_\pi} \\ &= \frac{-i}{(2\pi^4)} \int d^4p Tr(\gamma_0 S) \end{aligned}$$

S : nucleon propagator

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 41 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



The chemical potential is introduced in Euclidean space, p_0 is imaginary.


The integral has the form

$$\int d^4 p = \int d^3 \vec{p} \int_{-i\infty}^{i\infty} dp_0$$

When $\mu = 0$ the baryon number density vanishes, so

$$n(\mu_b) = \frac{-i}{(2\pi^4)} \int d^3 \vec{p} \left(\int_{-i\infty + \mu_b}^{i\infty + \mu_b} dp'_0 - \int_{-i\infty}^{i\infty} dp'_0 \right) \text{Tr} \left(\frac{\gamma_0}{\not{p}' - m - \Sigma(p')} \right)$$

the integral path in the complex p_0 plane:



[访问主页](#)

[标 题 页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 42 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



The usual χPT requires real momentum. It is unreasonable to extend the results to the complex plane when singularity appears according to perturbation theory. So nonperturbative method is necessary. Here we employ the same method as that in

J.A. Oller and E. Oset, Nucl. Phys. A 620, 438 (1997)

J.A. Oller, E. Oset and J.R. Pelaez, Phys. Rev. Lett. 80:3452 (1998)

to calculate the nucleon propagator.

The full nucleon propagator satisfy the following DSE:

$$S = S_0 + S_0 \Sigma S$$


S_0 : free nucleon propagator. Σ : self-energy

Replacing the full nucleon propagator in the self-energy by the free one, one obtains

$$S = (1 - S_0 \Sigma)^{-1} S_0 = (S_0^{-1} (1 - S_0 \Sigma))^{-1} = \frac{1}{S_0^{-1} - \Sigma}$$

Σ can be calculated perturbatively with the chiral effective Lagrangian.

the one loop diagrams contributing to Σ to $O(p^4)$ order:



访问主页

标题页

◀ ▶

◀ ▶

第 43 页 共 49 页

返 回

全屏显示

关 闭

退 出



explicit expression of the self-energy:

$$\begin{aligned}\Sigma &= -4c_1M^2 + \Sigma_a + \Sigma_b + \Sigma_c + e_1M^4 + O(p^5), \\ \Sigma_a &= \frac{3g_A^2}{4F_\pi^2}(m + \not{p}')\{M_\pi^2I + (m - \not{p}')\not{p}'I^{(1)}\}, \\ \Sigma_b &= \frac{3M_\pi^2\Delta_\pi}{F_\pi^2}\{2c_1 - \frac{p'^2}{m^2d}c_2 - c_3\}, \\ \Sigma_c &= -4c_1M_\pi^2\frac{\partial\Sigma_a}{\partial m}\end{aligned}$$

where $d = 4$, $\not{p}' = \not{p} + \gamma_0\mu_b$.

$$\begin{aligned}I &= -\frac{1}{8\pi^2}\frac{\alpha\sqrt{1-\Omega^2}}{1+2\alpha\Omega+\alpha^2}ArcCos\left(-\frac{\Omega+\alpha}{\sqrt{1+2\alpha\Omega+\alpha^2}}\right) - \frac{1}{16\pi^2}\frac{\alpha(\Omega+\alpha)}{1+2\alpha\Omega+\alpha^2}(2ln\alpha - 1), \\ I^{(1)} &= \frac{1}{2p'^2}\{(p'^2 - m^2 + M_\pi^2)I + \Delta_\pi\}, \\ \Delta_\pi &= \frac{M_\pi^2}{8\pi^2}ln\frac{M_\pi}{m}\end{aligned}$$

$$\begin{aligned}\alpha &= \frac{M_\pi}{m}, \\ \Omega &= \frac{p'^2 - m^2 - M_\pi^2}{2mM_\pi}\end{aligned}$$

The renormalization scheme we employed is the one based on "Infrared Regularization" proposed by Leutwyler et al. (Euro. Phys. J. C 9, 643 (1999)).

[访问主页](#)

[标题页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 44 页 共 49 页

[返 回](#)

[全屏显示](#)


[关 闭](#)

[退 出](#)

baryon number density:

$$n(\mu_b) = \frac{-i}{(2\pi^4)} \int d^4 p Tr \left(\frac{\gamma_0}{\not{p}' - m - \Sigma(\not{p}')} \right)$$

The integrand has three poles and one branch cut in the complex p_0 plane.



Position of poles: $p_0 = \sqrt{\vec{p}^2 + p_n^2}$, $n = a, b, c$. $p_a = 938$, $p_b = 1152 + i337$, $p_c = 1152 - i337$
(unit: MeV)

The branch cut starts at d, where $p_0 = \sqrt{\vec{p}^2 + (m + M_\pi)^2}$.

[访问主页](#)

[标 题 页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 45 页 共 49 页

[返 回](#)

[全 屏 显 示](#)


[关 闭](#)

[退 出](#)

baryon number susceptibility:

$$\chi(\mu_b) = \frac{\partial}{\partial \mu_b} n(\mu_b)$$

results for baryon number density and baryon number susceptibility:



critical chemical potential: $\mu = m_N = 938$ MeV

Discussion: Physically, after considering the binding energy, the value of the critical chemical potential will be less than 938 MeV. In our calculation the pion diagrams are not considered, so we have not revealed this contribution. It is expected that when calculating to higher orders, the result will be better.

When μ_b starts to be larger than 938 MeV, our result for the baryon number density is larger than the free nucleon result. When μ_b reaches 1152 MeV, the rate of the increasing of $n(\mu_b)$ begins to slow down. This value coincides with the chemical potential at which the $\chi(\mu_b) - \mu_b$ curve shows a peak.



[访问主页](#)

[标 题 页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 46 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)


[退 出](#)



EOS

$$n(\mu_b) = \frac{\partial P(\mu_b)}{\partial \mu_b}$$

$P - n$ curve



Our results on the baryon number density and EOS are very similar to the free nucleon approximation results. We think this explains why one can adopt the almost independent particle model in the study of nuclear matter.

[访问主页](#)

[标题页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 47 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)



★ Summary

- We give a direct method for calculating the EOS of QCD at zero T and finite μ
- We calculate the EOS of QCD at zero T and finite μ using four different QCD methods/models (rainbow-ladder DSE, HDL approximation, quasi-particle model, a nonperturbative approach inspired by χPT)

[访问主页](#)

[标 题 页](#)

[◀](#) [▶](#)

[◀](#) [▶](#)

第 48 页 共 49 页

[返 回](#)

[全 屏 显 示](#)

[关 闭](#)

[退 出](#)



[访问主页](#)

[标题页](#)

[◀◀](#) [▶▶](#)

[◀](#) [▶](#)

第 49 页 共 49 页

[返 回](#)

[全屏显示](#)

[关 闭](#)

[退 出](#)

Thank You!